



## *Botox / Dysport Treatment Before & After FAQs*

### ***Preparing for your treatment...***

5 DAYS BEFORE treatment discontinue the use of blood thinners like aspirin, Advil, Aleve, and fish oils as these can increase the risk of bruising & swelling after your *Botox/Dysport* treatment. Tylenol is preferred if over-the-counter pain reliever is needed.

Do not drink alcoholic beverages 24 hours before (and after) your treatment to avoid extra bruising.



***Do not use Botox/Dysport if you are pregnant or breastfeeding, are allergic to any of its ingredients, or suffer from neurological disorders. Please speak with the physician prior to your injection if you have any questions about any of these disqualifying conditions.***

*Failure to comply with recommendations & instructions  
may result in adverse effectiveness of treatment/therapy and may  
cause or worsen the occurrence of side effects*

***Common side effects: Injection site irritation & skin changes (itching, burning, bruising, dryness), headache, and sometimes dry/itchy eyes.***

***Contact our office if you experience: difficulty swallowing/breathing/talking, eye & vision problems, nerve/ muscle/ movement issues. Most side effects appear within one week of Botox/Dysport injection.***

TSC Aesthetics • 419 S. 5<sup>th</sup> Street, Gadsden, AL • 256-547-5200  
The Surgery Clinic, LLC After-Hours Emergency Line: 256-680-7035